

Elliptic flow fluctuations in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV

P. Sorensen^a for the STAR Collaboration*

^aBrookhaven National Laboratory, Upton, New York 11973-5000, USA

Please note that after these results were reported at Quark Matter 2006 and posted on the preprint server it was found that what is reported here as *elliptic flow fluctuations*, should rather be taken as an upper limit on the fluctuations. Further analysis has shown that fitting the multiplicity dependence of the q-distribution does not enable one to disentangle non-flow and fluctuations. The data from the q-vector distribution does not, therefore, exclude the case of zero fluctuations. The remainder of these proceedings we leave as they were originally reported.

We report first measurements of elliptic flow (v_2) fluctuations for the STAR collaboration at middle rapidity in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions. We analyze the multiplicity dependence of the flow vector length distribution to disentangle non-flow correlations from v_2 fluctuations. We find that the width of the v_2 distribution is approximately 36% of the mean v_2 and, within errors, independent of collision centrality. This value coincides with eccentricity fluctuations, apparently leaving little room for other sources of fluctuations.

Introduction: A primary goal of the RHIC program is to collide heavy-ions to create a quark-gluon plasma (QGP) and study its properties. Measurements of the azimuthal anisotropy of particle production with respect to the reaction plane (*i.e.* v_2) [1] seem to indicate that the matter created in collisions at RHIC behaves as a perfect fluid with a viscosity near a conjectured lower bound [2]. This conclusion is based primarily on comparisons to hydrodynamic model predictions [3, 1]. Uncertainty about the conditions at the beginning of the hydrodynamic expansion, however, make the conclusion that the matter created at RHIC has a viscosity near the conjectured lower bound ambiguous [4]. Since v_2 reflects the initial spatial eccentricity of the overlap region when two nuclei collide, fluctuations of v_2 should reflect fluctuations in the initial eccentricity. As such, v_2 fluctuations may provide sensitivity to the initial conditions and help determine the viscosity and other properties of the matter created in heavy-ion collisions.

Analysis method: The n^{th} harmonic reduced flow vector is defined as $q_{n,x} = \frac{1}{\sqrt{M}} \sum_i \cos(n\phi_i)$ and $q_{n,y} = \frac{1}{\sqrt{M}} \sum_i \sin(n\phi_i)$, where M is the number of tracks and ϕ_i is the azimuth angle of a track with respect to the reaction plane [5]. When enough tracks are used in the calculation of q_n , the central-limit-theorem (CLT) ensures that its distribution will be a two-dimensional Gaussian shifted by $\sqrt{M}v_n$ in the x direction with widths:

$$\sigma_x^2 = \frac{1}{2} (1 + v_{2n} - 2v_n^2 + g_n), \quad \sigma_y^2 = \frac{1}{2} (1 - v_{2n} + g_n). \quad (1)$$

*For the full list of STAR authors and acknowledgements, see appendix ‘Collaborations’ of this volume

g_n represents the broadening of the distribution that arises from non-flow correlations [5]. The exact direction of the reaction plane is not known, so we calculate the magnitude of the flow vector $|q_n|$. If v_2 does not fluctuate, the distribution of $|q|$ should be given by:

$$\frac{dN}{q_n dq_n} = \frac{1}{\sqrt{\pi}\sigma_x\sigma_y} e^{-\frac{1}{2}\left(\frac{q_n^2 + Mv_n^2}{\sigma_x^2}\right)} \sum_{k=0,2,4,\dots}^{\infty} \left(1 - \frac{\sigma_x^2}{\sigma_y^2}\right)^k \left(\frac{q_n}{v_n\sqrt{M}}\right)^k \frac{1}{k!} \Gamma\left(\frac{2k+1}{2}\right) I_k\left(\frac{q_n v_n \sqrt{M}}{\sigma_x^2}\right) \quad (2)$$

where Γ is the gamma function and I_k are modified Bessel's functions. Here we consider the $n = 2$ case. If v_2 fluctuates from event to event, the final $|q_2|$ distribution will have the form: $\frac{d\tilde{N}}{q_2 dq_2} = \int dv_2 f(v_2) \frac{dN}{q_2 dq_2}$, where $f(v_2)$ is the probability for an event to have a given v_2 . Zero v_2 fluctuations corresponds to $f = \delta(v_2 - \langle v_2 \rangle)$ where δ is the Dirac delta function. In our analysis we assume a Gaussian shape for the v_2 distribution: $f = \frac{1}{\sqrt{2\pi}\sigma_{v_2}} \exp\left(-\frac{(v_2 - \langle v_2 \rangle)^2}{2\sigma_{v_2}^2}\right)$, and extract the r.m.s. (σ_{v_2}) of the distribution from fits to data. Different assumed shapes change the r.m.s. by at most 15% (relative).




Figure 1. Simultaneous fits to three dN/dq distributions calculated using the same event class but using different fractions of the total multiplicity. The left panel shows a fit to simulated data and the right panel shows a fit to preliminary STAR data.

Non-flow and v_2 fluctuations both broaden the width of the observed q_2 distribution. This ambiguity is removed by using different numbers of tracks to calculate q_2 . Since the v_2 shift of the $|q_2|$ distribution scales with $\sqrt{M}$, the broadening of the distribution from v_2 fluctuations will scale with $\sqrt{M}$. We randomly discard tracks within an event class to calculate q_2 with reduced multiplicities. Then we fit the different $|q_2|$ distributions simultaneously to determine g_2 , σ_{v_2} , and $\langle v_2 \rangle$. Typical fits are shown in Fig. 1 for simulated data (left) and for real data (right). Best fit values agree well with the input parameters of the simulation. For our data analysis we use 15 million Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV measured with the STAR detector [6].

Eccentricity (standard and participant), impact parameter b , N_{bin} , N_{part} and multiplicity are calculated from a Monte-Carlo Glauber model [7] tuned to reproduce the multiplicity distribution of 200 GeV Au+Au collisions. We also use the model to subtract impact parameter fluctuations. The σ_{v_2} we present represents the width of the v_2 distribution at fixed b . The correction is carried out according to $\delta\sigma_{v_2} = \frac{\partial v_2}{\partial \varepsilon_{part}} \delta\sigma_{\varepsilon,part}$, where $\delta\sigma_{\varepsilon,part}$ is extracted from the difference between the width of the ε_{part} distribution when b is allowed to fluctuate and when it is fixed.

Results: The $\langle v_2 \rangle$ from our fits are shown versus ε_{part} in the top panel of Fig. 2 (left). Measuring v_2 fluctuations allows us to remove the major source of systematic uncertainty for $\langle v_2 \rangle$ [7, 9]. $\langle v_2 \rangle$ projects to zero for $\varepsilon_{part} = 0$ [8] giving us confidence that our measurements are sensitive to the participant axis (a coordinate system which can fluctuate to a different angle than the reaction plane [10]). We also note that $v_2 \not\propto \varepsilon_{part}$ as would be expected from hydrodynamic models [3]. The top panel of Fig. 2 (right) shows $\langle v_2 \rangle$ versus b compared to a phenomenological model $v_2 = 0.034\varepsilon_{part} (dN/dy)^{1/3}$ which is closely related to v_2/ε versus $\frac{1}{S}dN/dy$ plots [11]. Our data are consistent with this model.




Figure 2. The mean of the v_2 distribution ($\langle v_2 \rangle$) (top panels) and the r.m.s. width of the the distribution (σ_{v_2}) scaled by the mean (bottom panels). Data are presented versus participant eccentricity (left panel) and impact parameter (right panel). Various curves are explained in the text. Impact parameter fluctuations have been removed so that the data points represent the v_2 fluctuations for a fixed value of b .

The r.m.s. width over the mean of v_2 ($\sigma_{v_2}/\langle v_2 \rangle$) is shown in the bottom panels of Fig. 2. σ_{v_2} is corrected for b fluctuations so that the measurements represent the width of the distribution for a fixed impact parameter value. $\sigma_{v_2}/\langle v_2 \rangle$ is approximately 0.36 ± 0.03 and within errors independent of centrality. In the top left panel, we also show a comparison of the $\langle v_2 \rangle$ and σ_{v_2} values from this analysis to v_2 from 2- and 4-particle cumulant analyses ($v_2\{2\}$ and $v_2\{4\}$) [1, 7]. The curves derived from the measured σ_{v_2} and $\langle v_2 \rangle$ agree with the previously measured $v_2\{2\}$ and $v_2\{4\}$. The comparison shows that the differences between $v_2\{2\}$ and $v_2\{4\}$ can be explained by fluctuations without invoking non-flow suggesting that any effects from non-flow on the integrated v_2 are less important. This is not necessarily true, however, at higher p_T where the ratio of $v_2\{4\}/v_2\{2\}$ decreases with p_T . This decrease is likely related to an increase in non-flow correlations from hard processes at high p_T .

In the bottom right panel of Fig. 2 we compare $\sigma_{v_2}/\langle v_2 \rangle$ to nucleon Monte-Carlo Glauber calculations of $\sigma_\varepsilon/\langle \varepsilon \rangle$ where the eccentricity has been calculated with respect to the reaction plane (standard) or the participant axis (participant). The ε_{std} calculations can be excluded by data, demonstrating that our measurements are sensitive to the initial con-

ditions. The relative widths of the v_2 and ε_{part} distributions are consistent. This leaves little room for other sources of fluctuations and suggests that the conversion of ε to v_2 is identically efficient for every event. This conclusion is contradicted however by the variation of $\langle v_2 \rangle / \varepsilon$ with $(dN/dy)^{1/3}$. This contradiction may indicate that the fluctuations are overestimated in the Monte-Carlo Glauber model or that other sources of fluctuation are small compared to the eccentricity fluctuations. In principle, different treatments of the initial conditions can yield different predictions for the $\sigma_\varepsilon / \langle \varepsilon \rangle$. For example, calculations of the eccentricity based on Colored Glass Condensate (CGC) yield eccentricity values approximately 30% larger than a Glauber model [4]. If the geometric fluctuations in the CGC calculation are the same as for the Glauber model, then the ratio $\sigma_{\varepsilon,CGC} / \langle \varepsilon_{CGC} \rangle$ will be 30% smaller. This would allow room for other sources of fluctuations, consistent with the observation that $\langle v_2 \rangle / \varepsilon$ is not constant.

Hydrodynamic model calculations with fluctuating initial conditions [12] are also consistent with our data when a QGP EOS is used. The use of a resonance gas EOS seems to increase the ratio of the rms over the mean above the observed fluctuations. This indicates that although we have sought an observable that is sensitive to the initial conditions, we may learn about the later evolution as well and may even be sensitivity to the EOS.

Conclusions: We've disentangled non-flow and fluctuations and extracted σ_{v_2} and $\langle v_2 \rangle$ versus centrality for Au+Au collisions at 200 GeV. $\sigma_{v_2} / \langle v_2 \rangle$ is 0.36 ± 0.03 and within errors independent of centrality. These measurements significantly improve the accuracy of $\langle v_2 \rangle$, are consistent with 2- and 4-particle cumulant measurements and demonstrate sensitivity to the initial conditions. Future work will focus on the system-size dependence, energy dependence, and covariance of v_2 for different sub-events (*e.g.* partitioned according to pseudo-rapidity and p_T).

REFERENCES

1. K. H. Ackermann *et al.* [STAR Collaboration], Phys. Rev. Lett. **86**, 402 (2001); C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **89**, 132301 (2002); J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **92**, 052302 (2004); J. Adams *et al.* [STAR Collaboration], Phys. Rev. C **72**, 014904 (2005); J. Adams *et al.* [STAR Collaboration], Phys. Rev. Lett. **95**, 122301 (2005); K. Adcox *et al.* [PHENIX Collaboration], Nucl. Phys. A **757**, 184 (2005); J. Adams *et al.* [STAR Collaboration], Nucl. Phys. A **757**, 102 (2005).
2. P. Kovtun, D. T. Son and A. O. Starinets, Phys. Rev. Lett. **94**, 111601 (2005).
3. P. Huovinen, P. F. Kolb, U. W. Heinz, P. V. Ruuskanen and S. A. Voloshin, Phys. Lett. B **503**, 58 (2001); D. Teaney, Phys. Rev. C **68**, 034913 (2003).
4. T. Hirano, U. W. Heinz, D. Kharzeev, R. Lacey and Y. Nara, Phys. Lett. B **636**, 299 (2006); T. Lappi and R. Venugopalan, Phys. Rev. C **74**, 054905 (2006); H. J. Drescher and Y. Nara, arXiv:nucl-th/0611017.
5. J. Y. Ollitrault, arXiv:nucl-ex/9711003; A. M. Poskanzer and S. A. Voloshin, Phys. Rev. C **58**, 1671 (1998); N. Borghini, P. M. Dinh and J. Y. Ollitrault, Phys. Rev. C **63**, 054906 (2001).
6. STAR Collaboration, C. Adler *et al.*, Nucl. Instr. Meth. A **499**, 624 (2003).
7. M. Miller and R. Snellings, arXiv:nucl-ex/0312008.

8. But not necessarily for $b = 0$ (Fig. 2, top right).
9. X. l. J. Zhu, M. Bleicher and H. Stoecker, Phys. Rev. C **72**, 064911 (2005); R. Andrade, F. Grassi, Y. Hama, T. Kodama and O. J. Socolowski, arXiv:nucl-th/0608067.
10. S. Manly *et al.* [PHOBOS Collaboration], Nucl. Phys. A **774**, 523 (2006); R. S. Bhalerao and J. Y. Ollitrault, Phys. Lett. B **641**, 260 (2006).
11. C. Alt *et al.* [NA49 Collaboration], Phys. Rev. C **68**, 034903 (2003); S. A. Voloshin [STAR Collaboration], AIP Conf. Proc. **870**, 691 (2006).
12. T. Osada, C. E. Aguiar, Y. Hama and T. Kodama, arXiv:nucl-th/0102011; C. E. Aguiar, Y. Hama, T. Kodama and T. Osada, Nucl. Phys. A **698**, 639 (2002).